

## 英語論文

1. No evidence of deregulated patched-hedgehog signaling pathway in trichoblastomas and other tumors arising within nevus sebaceous.  
Takata M, Tojo M, Hatta N, Ohara K, Yamada M, Takehara K  
*J Invest Dermatol*, 117: 1666-70, 2001.
2. Sentinel lymph node biopsy in patients with extramammary Paget's disease.  
Hatta N, Morita R, Yamada M, Echigo T, Hirano T, Takehara K, Ichianagi K, Yokoyama K  
*Dermatol Surg*, 30: 1329-1234, 2004.
3. Management of squamous cell carcinoma in a patient with recessive-type epidermolysis bullosa dystrophica.  
Yamada M, Hatta N, Sogo K, Komura K, Hamaguchi Y, Takehara K  
*Dermatol Surg*, 30: 1424-9, 2004.
4. Localised vulva La gerhans cell histiocytosis.  
Ishigaki H, Hatta N, Yamada M, Orito H, Takehara K  
*Eur J Dermatol*, 14: 412-4, 2004.
5. Implications of popliteal lymph node detected by sentinel lymph node biopsy.  
Hatta N, Morita R, Yamada M, Takehara K, Ichianagi K, Yokoyama K  
*Dermatol Surg*, 31: 327-30, 2005.
6. Clinical significance of serum levels of soluble intercellular adhesion molecule-1 and soluble L-selectin in malignant melanoma.  
Yamada M, Yanaba K, Takehara K, Sato S  
*Arch Dermatol Res*, 297: 256-60, 2005.
7. Regulation of local and metastatic host-mediated anti-tumor mechanisms by L-selectin and intercellular adhesion molecule-1.  
Yamada M, Yanaba K, Hasegawa M, Matsushita Y, Horikawa M, Komura K, Matsushita T, Kawasuji A, Fujita T, Takehara K, Steeber DA, Tedder TF, Sato S  
*Clin Exp Immunol*, 143: 216-27, 2006.
8. Host-derived MCP-1 and MIP-1a regulate protective anti-tumor immunity to localized and metastatic B16 melanoma.

Nakasone Y, Fujimoto M, Matsushita T, Hamaguchi Y, Huu DL, Yanaba M, Sato S, Takehara K,  
Hasegawa M

*Am J Pathol*, 180: 365-74, 2012.